



## International Journal of Mosquito Research

ISSN: 2348-5906

CODEN: IJMRK2

IJMR 2023; 10(3): 01-07

© 2023 IJMR

[www.dipterajournal.com](http://www.dipterajournal.com)

Received: 01-01-2023

Accepted: 03-02-2023

**Maysa M Hegazy**

Department of Biology, Faculty  
of Science, Jazan University,  
Jazan 45142, Saudi Arabia

**Mohamed M Baz**

Department of Entomology,  
Faculty of Science, Benha  
University, Benha 13518, Egypt

# Efficacy of some plant wastes and cigarette filter residues as alternative agents for controlling *Culex pipiens* (Diptera: Culicidae)

**Maysa M Hegazy and Mohamed M Baz**

DOI: <https://doi.org/10.22271/23487941.2023.v10.i3a.671>

### Abstract

Mosquito-borne diseases are among the most common public health problems in the world, especially in developing countries. The idea of evaluating certain agricultural and industrial waste products of natural origin as larvicidal agents against mosquitoes is both good and unconventional. Four agricultural waste extracts (Apricot kernels, Lipton packet tea, date palm kernels, and burnt rice straw) and one industrial waste extract (cigarette filters) were evaluated as insecticides against 3<sup>rd</sup> the larval instar of *Culex pipiens* *In vitro* and their effect on egg laying and hatching rates. According to the current result, all plant and industrial waste extracts had larvicidal activity, and cigarette filters (100% mortality) and Lipton packet tea (97%) were most effective, followed by date palm kernels (90%), burnt rice straw (84%), and apricot kernel extracts (79%), whereas the LC<sub>50</sub> values were 528.14, 582.42, 815.5, 976, and 1080.50 ppm with 2000 ppm after 24 h, respectively. Data showed that cigarette filters and apricot kernel extracts were more attractive media for female oviposition, and cigarette filters and date palm kernel extracts had significant effects on the decreasing hatching rate (12.42% and 24.58%). Cigarette filters, Lipton packet tea, and date palm kernels can be used safely and efficiently to control *Cx. pipiens* larvae.

**Keywords:** Larvicidal activity, plant waste extracts, *Culex pipiens*, cigarette filters

### 1. Introduction

Infectious diseases spread by mosquitoes pose a serious threat to the world's population at the present time. There are many mosquito-borne diseases, such as malaria, dengue fever, yellow fever, filariasis, Japanese encephalitis, and Zika [1, 2]. Some reports indicate that more than 1 million people die each year as a result of mosquito-borne diseases worldwide, affecting more than 700 million people [3]. Therefore, control of these vector-borne is a critical step to take.

One of the main methods of controlling vector populations is the use of insecticides [4, 5]. It is considered the most effective solution for controlling these pests, but most mosquitoes have developed resistance to most groups of pesticides, which hinders the effectiveness of insecticide-based control measures and thus the spread of transmitted diseases.

As a result, it is imperative to find non-chemical methods to control a group of arthropods that act as vectors of human disease [6] in a safer, more natural, and sustainable way that does not affect the environment. Research continues, and there is growing interest in natural phytochemicals because they are safer and more efficient than synthetic insecticides. Therefore, we focus on identifying which plants, particularly those that have not been researched or worked on, such as plant waste, are expected to have higher concentrations of bioactive phytochemicals.

Plants are one of the gifts of nature that can treat a variety of human disorders [7, 8]. Also, plant extracts contain broadly effective antimicrobials along with their industrial and multi-industry scope [9-11]. Use some agricultural waste materials to reduce the cost of producing natural pesticides, antimicrobial agents, and other beneficial agricultural products [12-14].

Cigarette butts (CBs) are among the most common types of litter in the world [15] and result in approximately 4.5 trillion cigarettes being littered each year. CB consists of some tobacco fibers, ash, and about 30% filter, along with some chemical additives. Tobacco leaves naturally contain a wide range of compounds, including nicotine and sugar esters [16].

**Corresponding Author:****Mohamed M Baz**

Department of Entomology,  
Faculty of Science, Benha  
University, Benha 13518, Egypt

In addition to many other characteristics, such as cigarette size, tobacco rod weight, wrapping paper porosity, and filter type [17], cigarettes may contain approximately 10% by weight of additives known to increase toxicity [18]. The hope that CB wastes can be used in mosquito control is largely supported by the efficacy of nicotine and its derivatives against a variety of insect vectors, including mosquitoes [19, 20].

In this study, we aimed to evaluate the larvicidal activity of some plant residues, such as apricot kernels, Lipton-packet tea, date palm kernels, and burnt rice straw, with the effect of cigarette filter residues as alternative agents against *Culex pipiens* larvae of the West Nile virus vector. The effect of these materials on obstructing the female oviposition was also conducted.

## 2. Materials and methods

### 2.1 Plant materials and extraction

The Apricot kernels, Lipton packets of tea, and date palms were purchased from the supermarket (Fig. 1). The rice straw plants were obtained from rice fields in the North Delta, Egypt (Fig. 1). The air-dried Apricot and date palm kernels and Lipton packet teas were washed thoroughly in tap water and dried under indoor conditions at room temperature ( $27 \pm 2$  °C) for 11 days. The air-dried materials were powdered separately using a commercial electrical blender. For the extraction, the plant materials were extracted in water (aqueous extraction). Initially, the powder of 10 g of each plant waste was soaked in 200 ml of water for 48 h and filtered through Whatman filter paper No. 1. While the cigarettes consist of a filter for inhaling burnt tobacco (smoke) and a part of the tobacco wrapped inside a thin piece of paper to be the largest part of the cigarette (Fig. 1). In most cigarettes, specifically Marlboro cigarettes, after the smoking process, the cigarette is thrown out as cigarette butts with 1 to 2 cm of unburned tobacco and the cigarette butt filter. Cigarette butt filters were collected, weighed, and soaked in 100 ml of water for 48 h.

## 2.2 Mosquitocidal bioassay

### 2.2.1 Mosquito colony

*Culex pipiens* larvae were obtained from the Insect Breeding Laboratory, Department of Insects, Faculty of Science, Benha University, Egypt, and were used in all research experiments. Larvae were reared in tubs (25 x 20 x 10) filled with 2 L of dechlorinated tap water, Tetramin® fish meal, and dog biscuit powder every 2 days under laboratory conditions ( $27 \pm 2$  °C,


75–80% RH, and 12:12 h (L/D photoperiod). When *Cx. pipiens* pupae became adults, they were collected and placed in cages measuring 30x30x30 cm in insect cages, and then transferred to cups measuring 12x10x7 cm. An 8–10% sucrose solution was given as food to adults. After 48 hours of blood feeding, egg bowls filled with distilled water were placed in adult cages. Regularly, egg rafts were collected and placed in white enamel dishes filled with 1 to 1 liter of dechlorinated water. Larvae and adults were exposed to identical laboratory conditions [21].

### 2.2.2 Larvicidal activity

According to WHO [22], plant material extracts were investigated for their effectiveness against the third larvae of *Cx. pipiens*. The concentrations of plant waste materials and cigarette butt filters (125, 250, 500, 1000, 1500, and 2000 ppm) were performed [22]. The control groups were treated with water only. Twenty larvae were placed in a glass beaker holding 250 ml with various concentrations. Only water was used to treat the experiment and the control group. Five times, the experiments were run. Mortality rates for *Cx. pipiens* larvae was measured 24, 48 h after treatment (PT) at  $27 \pm 2$  °C and 80% relative humidity.

### 2.2.3 Oviposition inhibition

Twenty adult females and adult males were collected in wooden cages measuring 30x30x30 cm according to WHO protocol [22]. Adults were allowed to starve for one day after being fed with a 10% sugar solution for three days through the cotton pads. On the fifth day, a Syrian hamster was given to starving females for a blood meal. Plant waste extracts were applied inside the cage as follows: (1) 500 ppm, (2) 1000 ppm and (3) tap water (no concentration) in plastic cups that have 500 ml from the previous solutions. The Plant waste concentrations and tap water cups were placed in the corner of the cage for oviposition. Above the waterline, females lay rafts of eggs. The number of eggs in each egg raft was counted using stereomicroscopes. According to Hassan *et al.* [23], the eggs were divided into two groups: hatched and non-hatched. The following equation, which is based on the work of Sheikh [24], was used to calculate the hatching eggs. The percentage of eggs that hatch is equal to  $A/B \times 100$ , where A is the total number of eggs hatched and B is the total number of eggs deposited, as well as control substrate egg samples. The hatching of the eggs was observed 48 h after their placement on the surface of the cups.



**Fig 1:** The plant waste materials: apricot kernels (a), Lipton-packets tea (b), date palm kernels (c), rice straw burnt (d) and cigarette filters (e).

## 2.3 Statistical analysis

The study was carried out using IBM SPSS Statistics for Windows Version 23.0 (IBM Corp., Armonk, NY, USA). The significance of the percentages of larval mortality was assessed using one-way analysis of variance (ANOVA), and Duncan's multiple range test was employed as the post hoc test. Data from lethal concentrations (LC<sub>50</sub>) were analyzed using probit theory. The significant levels were set at  $p < 0.05$ .

## 3. Results

### 3.1 larvicidal activity of the plant waste extracts.

The larvicidal effects of plant waste extracts—apricot kernels, Lipton packet tea, date palm kernels, burnt rice straw, and cigarette filters were tested against 3<sup>rd</sup> larvae *Culex pipiens* indicating that mortalities were increased by increasing concentration and exposure time.

The mortality % of *Culex pipiens* after 24 h of exposure at 1500 and 2000 ppm of plant waste extracts; apricot kernels, Lipton-packets tea, date palm kernels, burnt rice straw, and cigarette filters were 59, 87, 71, 63, and 90% at 1500 ppm, and 79, 97, 90, 84, and 100% at 2000 ppm, 24 h, whereas the LC<sub>50</sub> values (50%, median lethal concentration) of these extracts were = 1080.50, 582.42, 815.5, 976.21 and 528.14 ppm, respectively (Table 1&2).

Data showed that all plant waste extracts had larvicidal activity and Lipton packet tea (97%) was most effective

followed by date palm kernels (90%), and burnt rice straw (84%) extracts, while apricot kernel (79%) extract were less effective. Also, the results showed that cigarette filter extract (100%) with LC<sub>50</sub> value = 528.14, 415.91 ppm was more effective than Lipton packet tea (97, 100%) with LC<sub>50</sub> value = 597.42, 470.91 ppm, at 2000 ppm, 24 and 48 h PT, respectively (Table 1&2 and Fig. 2).

### 3.1.1 Oviposition responses of female *Culex pipiens*

Data in Table 3 showed the effect of the five plant waste extracts on the oviposition, hatching, and non-hatching of female *Culex pipiens*. The rate of deposited eggs varied considerably according to apricot kernels, Lipton packet tea, date palm kernels, burnt rice straws, and cigarette filter extracts. The plant waste extracts were more attractive for oviposition, whereas the number of eggs laid was 818, 333, 350, 300, and 1355 eggs per 15 females at 1000 ppm concentration compared with untreated groups (1517, 1490, 1440, 1545, and 1580, respectively).

Data showed that cigarette filters and apricot kernel extracts were more attractive for female mosquitoes to lay eggs on. While both cigarette filters and date palm kernel extracts had significant effects on the decreasing hatching rate, the hatching rates reached 9.88% and 19%, respectively (Table 3).

**Table 1:** Efficacy of the plant waste extracts of apricot kernel, Lipton-packet tea, date palm kernel, rice straw, and cigarette filter on mortality of *Culex pipiens* after 24 and 48 exposures.


Time	Conc. (ppm)	Apricot kernel	Lipton packet tea	Date palm kernel	Burnt rice straw	Cigarette filters
24	0	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>
	125	6±1.87 <sup>fB</sup>	11±1.87 <sup>fA</sup>	8±1.22 <sup>fB</sup>	6±1.87 <sup>fB</sup>	13±2.00 <sup>fA</sup>
	250	10±1.58 <sup>eB</sup>	20±1.58 <sup>eA</sup>	12±2.00 <sup>eB</sup>	10±1.58 <sup>eB</sup>	21±1.87 <sup>eA</sup>
	500	25±1.58 <sup>dD</sup>	33±3.39 <sup>dB</sup>	28±3.00 <sup>dC</sup>	27±2.00 <sup>dCD</sup>	36±2.45 <sup>dA</sup>
	1000	37±2.55 <sup>cE</sup>	57±4.64 <sup>cB</sup>	43±4.06 <sup>cC</sup>	40±3.54 <sup>cD</sup>	65±2.24 <sup>cA</sup>
	1500	59±3.67 <sup>bE</sup>	87±2.55 <sup>bB</sup>	71±5.10 <sup>bC</sup>	63±3.39 <sup>bD</sup>	90±3.54 <sup>bA</sup>
	2000	79±4.00 <sup>aE</sup>	97±2.00 <sup>aB</sup>	90±3.06 <sup>aC</sup>	84±2.45 <sup>aD</sup>	100±0.00 <sup>aA</sup>
48	0	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>gA</sup>	0±0 <sup>fA</sup>
	125	10±3.54 <sup>fB</sup>	16±1.87 <sup>fA</sup>	12±2.55 <sup>fB</sup>	11±2.92 <sup>fB</sup>	18±1.22 <sup>eA</sup>
	250	14±2.92 <sup>eC</sup>	26±1.00 <sup>eA</sup>	17±2.55 <sup>eB</sup>	14±2.92 <sup>eC</sup>	28±2.00 <sup>dA</sup>
	500	29±2.92 <sup>dD</sup>	41±1.87 <sup>dB</sup>	34±1.87 <sup>dC</sup>	31±3.32 <sup>dD</sup>	44±3.67 <sup>cA</sup>
	1000	41±3.32 <sup>cE</sup>	67±5.39 <sup>cB</sup>	49±4.00 <sup>cC</sup>	45±3.16 <sup>cD</sup>	74±4.30 <sup>bA</sup>
	1500	63±4.64 <sup>bE</sup>	92±4.06 <sup>bB</sup>	75±4.74 <sup>bC</sup>	68±2.55 <sup>bD</sup>	98±2.00 <sup>aA</sup>
	2000	85±3.16 <sup>aC</sup>	100±0.00 <sup>aA</sup>	98±2.00 <sup>aA</sup>	90±2.74 <sup>aB</sup>	100±0.00 <sup>aA</sup>

a, b & c: There is no significant difference ( $P > 0.05$ ) between any two means, within the same column have the same superscript letter. A, B & C: There is no significant difference ( $P > 0.05$ ) between any two means, within the same row have the same superscript letter.

**Table 2:** Lethal time values of plant waste extracts against *Culex pipiens* after 24 and 48 exposures.

Time	Plant waste	LC <sub>50</sub> (low-Up.)	LC <sub>90</sub> (low-Up.)	LC <sub>95</sub> (low-Up.)	Slope± SE	X <sup>2</sup> (sig.)
24	Apricot kernel	1080.50 (805.28-1618.94)	4981.72 (3715.34-12405.76)	7669.05 (5565.02-22756.08)	1.939±0.16	10.578 (0.031)
	Lipton packet tea	582.42 (360.41-936.65)	2140.81 (1754.45-5412.76)	3074.01 (2668.70-9163.61)	2.312±0.16	25.150 (0.000)
	Date palm kernel	815.5 (560.77-1211.64)	3643.39 (2917.88-9215.71)	5466.36 (4420.94-16456.71)	2.062±0.16	15.688 (0.003)
	Burnt rice straw	976.27 (707.82-1445.62)	4064.44 (3156.78-9577.10)	6089.68 (4680.22-16867.85)	2.069±0.17	12.673 (0.01)
	Cigarette filters	528.14 (309.45-823.54)	1789.19 (1447.38-4405.40)	2528.54 (2182.00-7279.41)	2.418±0.13	27.456 (0.000)
48	Apricot kernel	916.47 (609.87-1547.48)	4636.69 (3816.19-15893.38)	7341.74 (6172.98-31979.08)	1.820±0.15	16.444 (0.002)
	Lipton packet tea	470.91 (268.60-732.58)	1707.27 (1368.02-4289.48)	2459.62 (2105.59-7296.54)	2.291±0.15	26.331 (0.000)
	Date palm kernel	645.34	2757.77	4109.33	2.097±0.15	31.744 (0.000)

		(361.61-1240.05)	(2621.19-11772.26)	(4422.55-23153.56)		
	Burnt rice straw	807.06 (506.28-1378.93)	3712.49 (3203.30-12797.70)	5721.89 (5196.98-25025.61)	1.933±0.15	20.653 (0.000)
	Cigarette filters	415.91 (245.02-684.97)	1728.66 (1399.85-4754.41)	2554.43 (2206.93-8559.86)	2.212±0.16	22.719 (0.000)



**Fig 2:** Larvicidal activity of plant waste materials; apricot kernels, Lipton packet tea, date palm kernels, burnt rice straw, and cigarette filters after 48 h of exposure.

**Table 3:** The impact of plant waste extracts on oviposition rate, egg hatching (%) and non-hatching eggs of *Culex pipiens*

Plant	Conc. (ppm)	Apricot kernel	Lipton packet tea	Date palm kernel	Burnt rice straw	Cigarette filters
Total egg	Control (0)	1517	1490	1440	1545	1580
	500	500	433	333	380	720
	1000	818	333	320	300	1355
No. of eggs hatched	Control (0)	1467	1432	1363	1477	1517
	500	427	333	243	282	333
	1000	370	152	207	182	277
No. of eggs non-hatched	Control (0)	50	58	77	68	63
	500	73	100	90	98	387
	1000	448	181	113	128	1078
Total hatching/ control (%)	Control (0)	100.00 <sup>aA</sup>	100.00 <sup>aA</sup>	100.00 <sup>aA</sup>	100.00 <sup>aA</sup>	100.00 <sup>aA</sup>
	500	29.09 <sup>bA</sup>	23.28 <sup>bAB</sup>	17.85 <sup>bB</sup>	19.07 <sup>bB</sup>	21.98 <sup>bB</sup>
	1000	25.23 <sup>bA</sup>	10.64 <sup>cC</sup>	15.16 <sup>bBC</sup>	12.30 <sup>cBC</sup>	18.24 <sup>bB</sup>
Total non-hatching/control (%)	Control (0)	0.00 <sup>bA</sup>	0.00 <sup>cA</sup>	0.00 <sup>bA</sup>	0.00 <sup>cA</sup>	0.00 <sup>bA</sup>
	500	70.91 <sup>aB</sup>	76.72 <sup>bAB</sup>	82.15 <sup>aA</sup>	80.93 <sup>bA</sup>	78.02 <sup>aA</sup>
	1000	74.77 <sup>aC</sup>	89.36 <sup>aA</sup>	84.84 <sup>aAB</sup>	87.70 <sup>aAB</sup>	81.76 <sup>aB</sup>
Hatching (%)	Control (0)	96.70 <sup>aA</sup>	96.09 <sup>aA</sup>	94.68 <sup>aA</sup>	95.58 <sup>aA</sup>	95.99 <sup>aA</sup>
	500	85.33 <sup>aA</sup>	76.92 <sup>bA</sup>	73.00 <sup>bA</sup>	74.12 <sup>bA</sup>	46.30 <sup>bB</sup>
	1000	45.21 <sup>bA</sup>	40.70 <sup>cA</sup>	24.58 <sup>cA</sup>	60.56 <sup>bA</sup>	12.42 <sup>cB</sup>
Non hatching (%)	Control (0)	3.30 <sup>bA</sup>	3.91 <sup>cA</sup>	5.32 <sup>bA</sup>	4.42 <sup>bA</sup>	4.01 <sup>cA</sup>
	500	14.67 <sup>bB</sup>	23.08 <sup>bB</sup>	27.00 <sup>aB</sup>	25.88 <sup>aB</sup>	53.70 <sup>bA</sup>
	1000	54.79 <sup>aB</sup>	54.30 <sup>aB</sup>	65.42 <sup>aB</sup>	39.44 <sup>aB</sup>	84.58 <sup>aA</sup>

a, b & c: There is no significant difference ( $P>0.05$ ) between any two means for each attribute, within the same column have the same superscript letter. A, B & C: There is no

significant difference ( $P>0.05$ ) between any two means for each attribute, within the same row have the same superscript letter.



#### 4. Discussion

Mosquito-borne diseases are among the most common public health problems in developing countries [25, 26]. It can be controlled by preventing mosquito bites by using repellents or hindering female mosquitoes from laying eggs, as well as by controlling mosquito larvae by safe methods, which leads to the death of many larvae and adult mosquitoes. Elimination of the larval stage is also beneficial because mosquito larvae cannot disperse or acquire human pathogens [27, 28].

The idea of evaluating some agricultural and industrial waste products of natural origin as larvicide agents against mosquitoes is a good and unconventional one since mosquito control requires new and innovative mechanisms that tends to nature. The primary criterion in selecting these compounds was the conversion of plant waste into useful materials, and they were easy and cheap to produce on a large scale.

Several previous studies were conducted mainly on botanical extracts of local plants in Egypt as well as around the world [28, 29-30] and their toxic effects on different types of insects [31].

In this study, the use of waste and its conversion into materials of value as an insecticide is evaluated. The tested extracts revealed differences in the types of plant and industrial waste extracts on the mortality rate of mosquito larvae, and these differences were attributed to the efficacy of the tested extracts and often to the main components of each.

Our findings demonstrated that cigarette filter extracts (100% mortality) were more effective than plant waste extracts at killing *Cx. pipiens* larvae, which had been exposed to all the plant waste extracts in our investigation after 24 and 48 h.

Several researchers recently evaluated cigarette butts, which are the unburned part of cigarette tobacco, along with a cigarette filter. The authors confirmed the toxicity of cigarette butts in killing mosquito larvae at a high rate, especially with increasing amounts of cigarette butts [32-35]. However, the effect of cigarette filters has not been studied; although they do not contain tobacco of any two flavors, burnt tobacco passes through them in addition to other components added to tobacco, which increases our belief that the cigarette filter contains toxic compounds that may reach or exceed the cigarette butts.

In a parallel study to our current work, Dieng *et al.* [36] recently studied the use of cigarette butts and their effect on changes in biological parameters from one generation to the next in mosquitoes. They examined the effects of three different concentrations of cigarette butt extract on mosquito egg-laying behaviors, egg hatchability, reproductive ability, longevity, and fertility. Finally, the mosquitoes that survived exposure to cigarette butts (CBs) waste had a short lifespan and altered life characteristics. As a result, the researchers confirmed the effectiveness of cigarette butt extract on mosquito larval mortality.

Previous literature showed that cigarette residues are huge wastes and amount to about 5.0 trillion [37] and that CBs contain toxic substances besides tobacco fibres, ash, and filtrate. US Department of Agriculture [38] referred to cigarettes as having about 10% by weight of additives known to increase toxicity [39].

Among the plant extracts, Lipton packet tea (97% mortality) and date palm kernel (90% mortality) were the most effective regarding lethal concentrations (LC<sub>50</sub>) and had lower values compared to other plant extracts at 24 and 48 h post-treatment, respectively.

It has been proven through similar studies that tea leaves

(*Camellia sinensis*) or green tea have an effective effect on killing mosquito larvae [28, 40-42]. The time of both larval and pupal development is significantly impacted when the young stages of the organism are exposed to *C. sinensis* leaf extract. These findings support a study by Dieng *et al.* [43] that indicated that dengue vectors treated to sublethal doses of *C. sinensis* leaf extracts had longer larval development times, lower pupation rates, and higher adult emergence rates. Because of its stimulating qualities, caffeine, a xanthine alkaloid, is a crucial component of green tea extract. Caffeine has been shown to interfere with the development of mosquito larvae [43] and may be effective against some pathogens [45].

We have hardly found studies on date palm kernels against mosquito larvae. Aqueous extracts of the fruit pulp, seed kernel, roots, bark, and leaves of *Balanites aegyptiaca* were tested for their ability to repel mosquito larvae of the *Cx. pipiens*. Larval mortality was highest in the aqueous root extract. In comparison to root and/or bark extracts, aqueous extracts of leaves, fruit pulp, and seed kernels revealed lower levels of larval mortality. All sections of *B. aegyptiaca* are thought to have larvicidal qualities that could be developed and utilised as organic pesticides to control mosquitoes [46].

Our data showed that the oviposition rate was higher in extracts of cigarette filters and apricot kernels, which were more attractive to female mosquitoes to lay eggs in. The reason for attracting female mosquitoes to lay eggs on the surface of extracted cigarette filters and apricot kernels is due to their light or dark brown color. Additionally, our data showed that palm kernel extracts and cigarette filters had a significant impact on the hatching rate. Where the percentage of hatching was low, which indicates the effect of these extracts on the death of mosquito egg embryos.

Our findings were consistent with those of Baeshen & Baz [47] and Zaitoun *et al.* [48], who investigated how *Acacia nilotica* extracts affected the ability to reduce egg hatchability and adult emergence. They discovered that acetone leaf extracts from *A. nilotica* were acutely toxic at 212.1 mg/L and chronically harmful at 144.2 mg/L.

Plant extracts contain a large, untapped stock of phytochemicals that could be widely used in mosquito control programs in place of synthetic insecticides. Several investigators have indicated the efficacy of phytochemicals against mosquito larvae based on their chemical nature and described the potential to kill mosquito larvae for a variety of plant-derived secondary substances [49].

Our findings concur with these findings. This study demonstrates that aqueous plant waste extracts of apricot kernels, Lipton packet tea, date palm kernels, burnt rice straw, and cigarette filters can be utilized as sustainable and eco-friendly mosquito repellents.

#### 5. Conclusion

It is necessary to stop the devastating effects of the illnesses that mosquitoes spread on people and their pets. The use of natural products as ecologically benign pesticides is currently a major area of research due to the wide variety and high potency of phytochemical substances. Thus, this work provided evidence of the effectiveness of plant waste (apricot kernels, Lipton packet tea, date palm kernels, and burnt rice straws) and one industrial waste extract (cigarette filters) as an eco-friendly and cost-effective insecticide against mosquito larvae. Therefore, we recommend the conversion and use of many agricultural residues in the control of mosquitoes and

insects of medical, veterinary, and economic importance. By transforming plant waste such as discarded rice straw into a useful, all-natural selective pesticide, the environmental issue posed by the rice straw can be resolved.

## 6. References

- World Health Organization (WHO). Dengue and severe dengue, accessed 10.08. 2013 <http://www.who.int/mediacentre/factsheets/fs117/en/>
- Benelli G, Jeffries, CL, Walker T. Biological control of mosquito vectors: Past, present, and future. *Insects*. 2016;7:52.
- Piovezan-Borges AC, Valente-Neto F, Urbietta GL, Laurence SGW, de Oliveira Roque F. Global trends in research on the effects of climate change on *Aedes aegypti*: International collaboration has increased, but some critical countries lag behind. *Parasit Vectors*. 2022;15:346.
- Van Den Berg H, Zaim M, Yadav RS, Soares A, Ameneshewa B, Mnzava A. Global trends in the use of insecticides to control vector-borne diseases. *Environmental health perspectives*. 2012;120(4):577-582.
- World Health Organization. Global insecticide use for vector-borne disease control: a 10-year assessment (2010–2019). World Health Organization. 2021.
- Pridgeon JW, Becnel JJ, Clark GG, Linthicum KJ. A high throughput screening method to identify potential pesticides for mosquito control. *J. Med. Entomol*. 2009;46:335-341.
- Deepa T, Elamathi R, Kavitha R, Kamalakannan SS, Suresh KJ. Screening for physical, phytochemical and antimicrobial activities of leaf extracts of *Sapindus emarginatus* Vahl. *Int. J Pharma. Tech. Res*. 2012;4(1):392-397.
- Baz MM, Khater HF, Baeshen RS, Selim A, Shaheen ES, El-Sayed YA, Salama SA, Hegazy MM. Novel Pesticidal efficacy of *Araucaria heterophylla* and *Commiphora molmol* extracts against camel and cattle blood-sucking ectoparasites. *Plants*. 2022;11(13):1682.
- Tembo Y, Mkindi AG, Mkenda PA, Mpumi N, Mwanauta R, Stevenson PC. Pesticidal plant extracts improve yield and reduce insect pests on legume crops without harming beneficial arthropods. *Frontiers in Plant Science*. 2018;9:1425.
- Hemeg HA, Moussa IM, Ibrahim S, Dawoud TM, Alhaji JH, Mubarak AS. Antimicrobial effect of different herbal plant extracts against different microbial population. *Saudi Journal of Biological Sciences*. 2020;27(12):3221-3227.
- Vieira DM, Pereira C, Calhella RC, Barros L, Petrovic J, Sokovic M, Machado AV. Evaluation of plant extracts as an efficient source of additives for active food packaging. *Food Frontiers*. 2022;3(3):480-488.
- Fahim AM, Kamel OM, Nawwar GAM. Facile synthesis of *in vitro* insecticidal and antimicrobial evaluation of bis heterocyclic moiety from PET waste recycling. *Egypt. Acad. J. Biolog. Sci.*, 2013;6(2):132-144.
- Eldiasty JG, Hassan MM, Kamel OM. Evaluation of some agricultural waste extracts against mosquito larvae, and some types of microorganisms as insecticidal and antibiotic agents. *Egyptian Academic Journal of Biological Sciences, G. Microbiology*. 2014;6(1):1-16.
- Farag SM, Essa EE, Alharbi SA, Alfarraj S, El-Hassan GA. Agro-waste derived compounds (flax and black seed peels): Toxicological effect against the West Nile virus vector, *Culex pipiens* L. with special reference to GC-MS analysis. *Saudi Journal of Biological Sciences*. 2021;28(9):5261-5267.
- Slaughter E, Gersberg RM, Watanabe K, Rudolph J, Stransky C, Novotny TE. Toxicity of cigarette butts, and their chemical components, to marine and freshwater fish. *Tob Control*, 2011 <http://dx.doi.org/10.1136/tc.2010.040170>.
- Leffingwell JC. Basic chemical constituents of tobacco leaf and differences among tobacco types. In: Davis, D.L., Nielsen, M.T. (Eds.), *Tobacco – Production, Chemistry and Technology*. Blackwell Science Ltd., London, 1999, 265–303.
- Perfetti TA, Coleman WM, Smith WS. Determination of mainstream and sidestream cigarette smoke components for cigarettes of different tobacco types and a set of reference cigarettes. *Beitr. Tabakforsch. Int*. 1998;18:95–113.
- Comité Scientifique des Risques Sanitaires Emergents et Nouveaux (CSRSE), 2010. Les additifs du tabac, accessed 16.08.2013 [http://ec.europa.eu/health/scientificcommittees/opinions\\_layman/tobacco/fr/about.htm#7](http://ec.europa.eu/health/scientificcommittees/opinions_layman/tobacco/fr/about.htm#7)
- Tomizawa M, Casida JE. Neonicotinoid insecticide toxicity: mechanisms of selective action. *Ann. Rev. Toxicol*. 2005;45:247–268.
- Sohail A, Hamid FS, Waheed A, Ahmed N, Aslam N, Zaman Q, Ahmed F, Islam S. Efficacy of different botanical materials against aphid *Toxoptera aurantii* on tea (*Camellia sinensis* L.) cuttings under high shade nursery. *J. Mater. Environ. Sci*. 2012;3:1065–1070
- Mohamed M Baz, Maysa M Hegazy, Hanem F Khater and Yasser A El-Sayed. Comparative Evaluation of Five Oil-Resin Plant Extracts against The Mosquito Larvae, *Culex pipiens* Say (Diptera: Culicidae). *Pakistan Veterinary Journal*. 2021;41(2):191-196.
- World Health Organization. Guidelines for Laboratory and Field Testing of Mosquito Larvicides; World Health Organization: Geneva, Switzerland, 2005.
- Hassan M, Zayed, A, Ahmed M. The influence of symbiotic bacteria on digestion and yolk protein synthesis in *Cx. pipiens* L. (Diptera: Culicidae). *Journal of Egyptian German Society of Zoology*. 1996;21(E):269-284.
- El-Sheikh T. Effects of application of selected insect growth regulators and plant extracts on some physiological aspects of the black cutworm, *Agrotis ipsilon*, (Huf). Ph.D. Thesis, Faculty of Science, Ain Shams University, 2002, 186.
- World Health Organization. Vector-Borne Diseases World Health Organization, 2017.
- Khan SA, Web CE, Kassim NFA. Prioritizing mosquito-borne diseases during and after the COVID-19 pandemic. *Western Pacific Surveillance and Response Journal: WPSAR*. 2021;12(2):40.
- Hardin JA, Jackson FLC. Applications of natural products in the control of mosquito-transmitted diseases. *Afr. J. Biotechnol*. 2009;8:7373–7378.
- Baz MM, Selim A, Radwan IT, Alkhaibari AM, Khater HF. Larvicidal and adulticidal effects of some Egyptian oils against *Culex pipiens*. *Scientific Reports*. 2022;12(1):1-18.

29. Khater H, Soliman E, Slim D, Debboun AM, Baz MM. Larvicidal Efficacy of Fifteen Plant Essential Oils against *Culex pipiens* L. Mosquitoes in Egypt. *Egyptian Journal of Veterinary Sciences*. 2023;54(2):183-192
30. Baz MM, Selim AM, Radwan IT, Khater HF. Plant oils in the fight against the West Nile Vector, *Culex pipiens*. *International Journal of Tropical Insect Science*, 2022, 1-8.
31. Kamel OM, Abd El-Baky SMM, Hassan MM, Hafez JA, Hamed MS. Synergistic and joint actions of some plant extracts on their larvicidal activity against the mosquito *Ochlerotatus caspius* (Diptera: Culicidae). *Egypt Acad Soc Environ Develop*. 2005;6(2):277-289.
32. Dieng H, Rajasaygar S, Ahmad AH, Ahmad H, Rawi CSM, Zuharah WF. Turning cigarette butt waste into an alternative control tool against an insecticide-resistant mosquito vector. *Acta tropica*. 2013;128(3):584-590.
33. Dieng H, Rajasaygar S, Ahmad AH, Rawi CSM, Ahmad H, Satho T. Indirect effects of cigarette butt waste on the dengue vector *Aedes aegypti* (Diptera: Culicidae). *Acta tropica*. 2014;130:123-130.
34. Mondal NK, Hajra A, Chakraborty D, Medda S, Dey U, Datta JK. Cigarette butt waste and its effective utilization towards larvicidal activity of mosquito. *International Journal of Scientific Research in Environmental Sciences*. 2015;3(1):9.
35. Panneerselvam C, Suresh U, Alatawi FA, Angayarkanni J, Murugan K. Cigarette butt waste derived activated carbon incorporated silver nanoparticle (AC-Ag): Effective nanocomposite for anti-bacterial and anti-larval activity in wastewater remediation. *Materials Letters*. 2022;313:131809.
36. Dieng H, Sudha R, Abu Hassan A, Hamdan A, Che Salmah MH, Wan Fatma Z. Turning cigarette butt waste into an alternative control tool against an insecticide-resistant mosquito vector. *Acta Trop*, 2013.
37. Slaughter E, Gersberg RM, Watanabe K, Rudolph J, Stransky C, Novotny TE. Toxicity of cigarette butts, and their chemical components, to marine and freshwater fish. *Tob Control*, 2011. <http://dx.doi.org/10.1136/tc.2010.040170>.
38. US Department of Agriculture (USDA), 2010. Tobacco statistics and reports, accessed 10.09.2013. <http://www.fas.usda.gov/cots/tobstats.html>
39. Wertz MS, Kyriass T, Paranjape S, Glantz SA. The toxic effects of cigarette additives. Philip Morris' project mix reconsidered: an analysis of documents released through litigation. *PLoS Med*. 2011;8(12):e1001145.
40. Das YK, Ghar SB. The implicit remedy to eradicate *Culex pipiens* with eco-friendly biointegrated lemongrass palladium metallic nanoparticles. *Int J Entomol Res*. 2022;7:8-23.
41. Hassan MI, Atwa WA, Moselhy WA, Mahmoud DA. Efficacy of the green tea, *Camellia sinensis* leaves extract on some biological activities of *Culex pipiens* and the detection of its phytochemical constituents. *Egyptian Academic Journal of Biological Sciences, F. Toxicology & Pest Control*. 2020;12(1):59-70.
42. Som DK, Iaha P, Mondal S. Larvicidal Effect of *Coffea arabica* L., *Camellia sinensis* (L.) Kuntz, and *Punica granatum* L. On *Aedes albopictus* (Skuse), The Vector of Dengue and Chikungunya. *Uttar Pradesh Journal of Zoology*, 2021, 10-18.
43. Dieng H, Tan Yusop NSB, Kamal NNB, Ahmad AH, Ghani IA. Exposure of a dengue vector to tea and its waste: Survival, developmental consequences, and significance for pest management. *J Agric Food Chem*. 2016;64:3485-3491.
44. Laranja AT, Manzatto AJ, Campos Bicudo HEM. Effects of caffeine and used coffee grounds on biological features of *Aedes aegypti* (Diptera, Culicidae) and their possible use in alternative control. *Genet. Mol. Biol*. 2003;26(4):419-29.
45. Van Breda SV, Van der Merwe CF, Robbertse H, Apostolides Z. Immunohistochemical localization of caffeine in young *Camellia sinensis* (L.) O. Kuntze (tea) leaves. *Planta*, 2012. DOI: 10.1007/s00425-012-1804-x.
46. Chapagain B, Wiesman Z. Larvicidal effects of aqueous extracts of *Balanites aegyptiaca* (desert date) against the larvae of *Culex pipiens* mosquitoes. *African journal of Biotechnology*, 2005, 4(11).
47. Baeshen RS, Baz MM. Efficacy of *Acacia nilotica*, *Eucalyptus camaldulensis*, and *Salix safsafs* on the mortality and development of two vector-borne mosquito species, *Culex pipiens* and *Aedes aegypti*, in the laboratory and field. *Heliyon*, 2023, 9(5).
48. Zaitoun AA, Madkour MH, Shamy MY. Effect of three plants extracts on some bacterial strains and *Culex pipiens* L. stages, *J Eng. Sci. Technol*. 2012;5:54-63
49. Chan YB, Selvanathan V, Tey LH, Akhtaruzzaman M, Anur FH, Djearmane S. Effect of Calcination Temperature on Structural, Morphological and Optical Properties of Copper Oxide Nanostructures Derived from *Garcinia mangostana* L. Leaf Extract. *Nanomaterials*. 2022;12(20):3589.
50. Muema JM, Bargul JL, Nyanjom SG, Mutunga JM, Njeru SN. Potential of *Camellia sinensis* proanthocyanidins-rich fraction for controlling malaria mosquito populations through disruption of larval development. *Parasites & vectors*. 2016;9(1):1-10.